

Day-Length During Seed Development Affects Size, Germinability and Storability of Lettuce Seeds.

S. Contreras¹, M. A. Bennett¹ and D. Tay²

¹*Dept. of Horticulture and Crop Science, Ohio State Univ., Columbus OH 43210-1086, USA*

²*Ornamental Plant Germplasm Center, Ohio State Univ., Columbus OH 43210-1086, USA*

Seed germinability and storability are important aspects of seed quality determined by the genotype and environment of seed development. The objective of this study was to determine how day-length of the mother plant environment affects lettuce seed quality. Seeds of cv. Tango were produced in growth chambers under one of two treatments: i) short day (SD), consisting of 8 h of fluorescent light ($\sim 310 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) plus 16 h of darkness daily, and ii) long day (LD), consisting of 8 h of fluorescent light plus 8 h of incandescent light ($\sim 21 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and 8 h of darkness daily. The LD treatment produced significantly heavier seeds, however germination at optimal conditions (20°C-light) was similar for both treatments. At suboptimal conditions (30°C, 20°C with different external ABA concentrations, or dark), seed germinability (% and rates) from SD treatment was higher. After accelerated aging (41°C, $\sim 100\%$ RH, 72 h) germination of normal seedlings was higher for seeds from LD. Seed germination was also evaluated after 2, 4, and 6 months of storage at 30°C, 74% RH. Stored seed presented a progressive and significant reduction of germinability for both treatments, however seeds from SD were more affected. The results indicated that day-length during seed development affected lettuce seed weight, germinability, and storability. In this case, germinability and storability were inversely related. The critical moment for day-length effects was also studied. Seed size showed to be determined earlier during seed development (first 6 days). Conversely, seed germinability and storability were determined at the end of seed development, after physiological maturity, which occurred ~ 11 days after flowering.